HSC Research Report

HSC/19/04

Regularized Quantile Regression Averaging for probabilistic electricity price forecasting

Bartosz Uniejewski¹
Rafał Weron¹

¹ Department of Operations Research, Faculty of Computer
Science and Management, Wrocław University of Science
and Technology, Poland

Hugo Steinhaus Center
Wrocław University of Science and Technology
Wyb. Wyspiańskiego 27, 50-370 Wrocław, Poland
<http://www.im.pwr.wroc.pl/~hugo/>

Regularized Quantile Regression Averaging for probabilistic electricity price forecasting

Bartosz Uniejewski^a, Rafał Weron^a

^a*Department of Operations Research, Wrocław University of Science and Technology, 50-370 Wrocław, Poland*

Abstract

Quantile Regression Averaging (QRA) has sparked interest in the electricity price forecasting community after its unprecedented success in the Global Energy Forecasting Competition 2014, where the top two winning teams in the price track used variants of QRA. However, recent studies have reported the method's vulnerability to low quality predictors when the set of regressors is larger than just a few. To address this issue, we consider a regularized variant of QRA, which utilizes the Least Absolute Shrinkage and Selection Operator (LASSO) to automatically select the relevant regressors. We evaluate the introduced technique – dubbed LASSO QRA or LQRA for short – using datasets from the Polish and Nordic power markets, a set of 25 point forecasts obtained for calibration windows of different lengths and 20 different values of the regularization parameter. By comparing against nearly 30 benchmarks, we provide evidence for its superior predictive performance in terms of the Kupiec test, the pinball score and the test for conditional predictive accuracy.

Keywords: electricity price forecasting, probabilistic forecast, Quantile Regression Averaging, LASSO, Kupiec test, pinball score, conditional predictive accuracy

1. Introduction

The Global Energy Forecasting Competition 2014 (GEFCom2014) has changed the landscape of energy forecasting. Instead of focusing on point forecasts, with all their limitations, the organizers have advocated computing *probabilistic* – more precisely – *quantile forecasts* instead and using the pinball score to evaluate them (Hong et al., 2016). The *electricity price forecasting* (EPF) literature has reacted lively, with the number of probabilistic EPF papers growing steadily in the following years, see Nowotarski and Weron (2018) for a recent review.

Formally introduced by Nowotarski and Weron (2015), *Quantile Regression Averaging* (QRA) has sparked interest in the EPF community after its unprecedented success in GEFCom2014, where the top two winning teams in the price track – Gaillard et al. (2016) and Maciejowska and Nowotarski (2016) – used variants of QRA. Since the method is a general forecasting technique not limited to electricity prices, a number of authors have reported its successful application

Email addresses: bartosz.uniejewski@pwr.edu.pl (Bartosz Uniejewski), rafal.weron@pwr.edu.pl (Rafał Weron)

in areas ranging from load (Liu et al., 2017; Zhang et al., 2018; Wang et al., 2019) to wind power (Zhang et al., 2016) and irradiance forecasting (Mpfumali et al., 2019).

However, a recent study of Marcjasz et al. (2020) has revealed the method’s vulnerability to low quality predictors when the set of regressors is larger than just a few. The authors emphasize the importance of correctly selecting the individual point forecasts for QRA, whose performance is known only *ex post*, and even recommend using only two or three point forecasts as inputs. A way to tackle this problem via principal component analysis (PCA) has been proposed by Maciejowska et al. (2016). However, their Factor QRA (or FQRA) model yields suboptimal and nearly identical forecasts to the so-called Quantile Regression Machine (QRM) of Uniejewski et al. (2019). Here, we introduce a different approach that outperforms not only these two, but also a number of other benchmarks. It can be regarded as a regularized variant of QRA, which utilizes the *Least Absolute Shrinkage and Selection Operator* (LASSO) of Tibshirani (1996) to select the relevant regressors. We call the method LASSO QRA or LQRA for short.

We evaluate the introduced technique using datasets from the Polish and Nordic power markets, a set of 25 point forecasts obtained for calibration windows of different lengths and 20 different values of the regularization parameter. Then we compare LQRA against nearly 30 benchmarks and provide evidence for its superior predictive performance in terms of the Kupiec (1995) test, the pinball score (see, e.g., Gneiting, 2011; Nowotarski and Weron, 2018) and the test for conditional predictive accuracy of Giacomini and White (2006).

The remainder of the paper is structured as follows. In Section 2 we briefly describe the datasets, then in Section 3 we discuss the rolling window scheme used in this study, the construction of point forecasts and introduce LQRA. In Section 4, we first evaluate the predictive performance of point forecasts, then discuss coverage and sharpness of the obtained probabilistic forecasts. Finally, in Section 5 we wrap up the results and conclude.

2. Datasets

We consider datasets from two European power markets – the Polish Power Exchange (POLEX) and Nord Pool (NP). The first one was downloaded from the websites of POLEX (www.tge.pl) and the Polish Transmission System Operator (TSO; www.pse.pl). It comprises three time series at hourly resolution (see Figure 1):

- day-ahead electricity prices for the main auction at POLEX (so-called ‘Fixing #1’),
- day-ahead predictions of the system-wide load in Poland,
- day-ahead predictions of the generation of centrally dispatched generating units in Poland (denoted by JWCD),

and covers the period from 2 October 2014 to 31 December 2018. The second dataset was downloaded from the Nord Pool website (www.nordpoolgroup.com/historical-market-data/) and also includes three time series at hourly resolution (see Figure 2):

- day-ahead system prices for the whole Nordic region,

Figure 1: Polish Power Exchange (POLEX) day-ahead prices (*top*), day-ahead system load forecasts (*middle*) and day-ahead forecasts of the generation of centrally dispatched generating units (*bottom*) from 2 October 2014 to 31 December 2019. The vertical dashed lines mark respectively the end of the initial 728-day calibration window for point forecasts and the end of the initial 364-day calibration window for probabilistic forecasts. The remaining observations constitute the 460-day out-of-sample test period.

- day-ahead load forecasts (so-called *consumption prognosis*) aggregated for Denmark, Finland, Norway and Sweden,
- day-ahead forecasts of wind power generation in Denmark,

and covers the same period from 2 October 2014 to 31 December 2018. All time series were pre-processed to account for changes to/from the daylight saving time. The missing values (corresponding to changes to the summertime) were substituted by the arithmetic average of the observations from neighboring hours. The doubled values (corresponding to the changes from the summertime) were replaced by their arithmetic mean.

3. Methodology

3.1. Calibration windows

Like the majority of EPF studies, we use a rolling window scheme. However, following Hubicka et al. (2019) and Marcjasz et al. (2018), we do not consider an arbitrarily chosen calibration

Figure 2: Nord Pool (NP) system prices (*top*), day-ahead consumption prognosis (*middle*) and day-ahead forecasts of wind power generation (*bottom*) from 2 October 2014 to 31 December 2019. The vertical dashed lines mark respectively the end of the initial 728-day calibration window for point forecasts and the end of the initial 364-day calibration window for probabilistic forecasts. The remaining observations constitute the 460-day out-of-sample test period.

window length, but rather a pool of 701 window lengths – ranging from 28 (ca. one month) to 728 days (ca. two years) – and combine the obtained forecasts. Initially, the first 728 ($= 2 \times 364$) days are used for model calibration. For shorter windows, the calibration sample is left-truncated, so that it ends on the same day as the 728-day window. For each day in the period 29 September 2016 to 31 December 2018, we simultaneously compute 24 point forecasts, one for each hour of the day. Furthermore, we use a rolling calibration window of 364 days (ca. one year) to obtain probabilistic forecasts for each hour of the days between 28 September 2017 and 31 December 2018. The forecasts for all 460 days of the out-of-sample period are evaluated in Section 4.2. Dashed lines in Figures 1 and 2 mark respectively the end of the initial 728-day calibration window for point forecasts (i.e., 2 October 2014 to 28 September 2016) and the end of the initial 364-day calibration window for probabilistic forecasts (i.e., 29 September 2016 to 27 September 2017).

3.2. Point forecasts

3.2.1. Variance stabilizing transformation

Before calibrating any models, we transform the data using the *area hyperbolic sine* (asinh) variance stabilizing transformation introduced by Uniejewski et al. (2018). Namely, each time series (prices and exogenous variables) is first standardized by subtracting the sample median and dividing by the sample *Median Absolute Deviation* (MAD) corrected by the 75% quantile of the standard normal distribution $z_{0.75}$. For instance, for prices $P_{d,h}$ we have:

$$p_{d,h} = \frac{P_{d,h} - a}{b} \equiv \frac{P_{d,h} - \text{med}(P_{d,h})}{\frac{\text{MAD}(P_{d,h})}{z_{0.75}}}.$$

Then we asinh-transform the standardized series. For instance, for prices we obtain:

$$Y_{d,h} = \text{asinh}(p_{d,h}) = \log\left(p_{d,h} + \sqrt{p_{d,h}^2 + 1}\right), \quad (1)$$

for each d and h . Once the transformed price forecast for day d and hour h , i.e., $\hat{Y}_{d,h}$, is computed, we apply the inverse transformation to obtain the price forecast itself. Here, we follow Narajewski and Ziel (2019) and define the back-transformation as:

$$\hat{P}_{d,h} = a + \frac{b}{D} \sum_{i=1}^D \sinh\left(\hat{Y}_{d,h} + \varepsilon_i\right), \quad (2)$$

where D is the number of days in the calibration window and ε_i are the in-sample residuals of the OLS procedure. Note, that the back-transformation proposed by Uniejewski et al. (2018) is simpler, but may be less accurate when $\hat{Y}_{d,h}$ is far from zero.

3.2.2. The expert model

For computing point forecasts we use a parsimonious autoregressive structure, inspired by the ARX model of Misiorek et al. (2006) and later used in a number of EPF studies (see Ziel and Weron, 2018, for a review). Following Uniejewski et al. (2016) and Ziel (2016), we refer to it as an *expert* model, since it is built on some prior knowledge of experts. The model for the asinh-transformed price on day d and hour h is given by:

$$\begin{aligned} Y_{d,h} = & \underbrace{\beta_1 Y_{d-1,h} + \beta_2 Y_{d-2,h} + \beta_3 Y_{d-7,h}}_{\text{autoregressive effects}} + \underbrace{\beta_4 Y_{d-1,24}}_{\text{end-of-day}} + \underbrace{\beta_5 Y_{d-1}^{\max} + \beta_6 Y_{d-1}^{\min}}_{\text{non-linear effects}} \\ & + \underbrace{\beta_7 Z_{d,h}^{(1)} + \beta_8 Z_{d,h}^{(2)}}_{\text{exogenous variables}} + \underbrace{\beta_9 D_{Sat} + \beta_{10} D_{Sun} + \beta_{11} D_{Mon}}_{\text{weekday dummies}} + \varepsilon_{d,h}, \end{aligned} \quad (3)$$

where $Y_{d-1,h}$, $Y_{d-2,h}$ and $Y_{d-7,h}$ account for the autoregressive effects and correspond to prices from the same hour of the previous day, two days before and a week before, $Y_{d-1,24}$ is the last known price at the time the prediction is made and provides information about the end-of-day price level, $Y_{d-1}^{\max} \equiv \max_{h=1,\dots,24}\{Y_{d-1,h}\}$ and $Y_{d-1}^{\min} \equiv \min_{h=1,\dots,24}\{Y_{d-1,h}\}$ represent previous day's variation in prices, $Z_{d,h}^{(1)}$ is the asinh-transformed day-ahead system load forecast for day d and hour h , $Z_{d,h}^{(2)}$ is

the asinh-transformed day-ahead forecast for day d and hour h of either generation of centrally dispatched generating units (for POLEX) or of wind power generation (for Nord Pool), D_{Sat} , D_{Sun} and D_{Mon} are respectively Saturday, Sunday and Monday dummies, and the $\varepsilon_{d,h}$'s are assumed to be independent and identically distributed normal variables. As is common in the EPF literature (see, e.g., Ziel and Weron, 2018), we consider 24 separate time series and calibrate the same model structure independently for each hour of the day. Finally, the model weights are estimated by minimizing the Residual Sum of Squares (RSS), independently for each calibration window length (ranging from 28 to 728 days, see Section 3.1).

3.3. Probabilistic forecasts

3.3.1. Quantile Regression Averaging (QRA)

QRA, as introduced by Nowotarski and Weron (2015), is based on averaging $n = 1, \dots, N$ point predictions of electricity prices $\hat{P}_{d,h}^{(n)}$ using quantile regression (QR; for a thorough account of QR see Koenker, 2005):

$$q(\alpha|X_{d,h}) = X_{d,h}\beta_\alpha, \quad (4)$$

where $q(\alpha|\cdot)$ is the conditional α th quantile of the electricity price distribution, $X_{d,h}$ is the matrix of independent variables, i.e., $\hat{P}_{d,h}^{(n)}$ for $n = 1, \dots, N$, while $\beta_\alpha = [\beta_\alpha^{(1)}, \dots, \beta_\alpha^{(N)}]'$ is the corresponding vector of weights for given α . To obtain parameter estimates we minimize the sum of the so-called *check functions*:

$$\hat{\beta}_\alpha = \underset{\beta_\alpha}{\operatorname{argmin}} \left\{ \sum_{d,h} \underbrace{\left(\alpha - \mathbb{1}_{P_{d,h} < X_{d,h}\beta_\alpha} \right) (P_{d,h} - X_{d,h}\beta_\alpha)}_{\text{check function}} \right\}. \quad (5)$$

Following the unprecedented success in the Global Energy Forecasting Competition 2014, QRA became a popular technique for probabilistic energy forecasting (see, e.g., Maciejowska et al., 2016; Zhang et al., 2016; Liu et al., 2017; Zhang et al., 2018; Kostrzewski and Kostrzewska, 2019; Mpfumali et al., 2019; Serafin et al., 2019; Uniejewski et al., 2019; Wang et al., 2019). However, a recent study of Marcjasz et al. (2020) has revealed the method's vulnerability to low quality predictors when the set of regressors is larger than just a few. To tackle this problem, in what follows we introduce a *regularized* (or *penalized*) variant of QRA.

QRA is a special case of QR where the independent variables (or regressors) are point forecasts of the dependent (or predicted) variable, i.e., $\hat{P}_{d,h}^{(n)}$ for $n = 1, \dots, N$. Note, however, that QR is a general technique that can yield electricity price forecasts also for more diverse sets of explanatory variables, including past prices and/or fundamental variables, as in Bunn et al. (2016) or Hagfors et al. (2016).

3.3.2. Regularization

In simple terms the idea of regularization can be formulated as an optimization problem:

$$\hat{\beta} = \underset{\beta}{\operatorname{argmin}} \{ f(X; \beta) + g(\beta) \}, \quad (6)$$

where $f(X; \beta)$ is the minimized function, $g(\beta)$ is the penalty function and β is the vector of parameters (Tikhonov, 1963). The most commonly used variant of regularization defines the penalty function as an ℓ^q norm scaled with an additional *tuning* (or *regularization*) parameter λ . When

$q = 1$ and $f(X; \beta)$ is the Residual Sum of Squares we obtain the *Least Absolute Shrinkage and Selection Operator* (LASSO) of Tibshirani (1996):

$$\hat{\beta} = \underset{\beta}{\operatorname{argmin}} \left\{ \text{RSS} + \lambda \sum_{i=1}^n |\beta_i| \right\}, \quad (7)$$

which not only shrinks the β 's towards zero, but also eliminates some of them. For $\lambda = 0$ the method is equivalent to the original formulation of the problem (i.e., Ordinary Least Squares, OLS). As λ increases, more and more variables are eliminated. A clear advantage of LASSO is that it can handle an almost unlimited number of explanatory variables. Hence, it can be used as an automated variable selection tool (Gaillard et al., 2016; Uniejewski et al., 2016; Ziel, 2016; Uniejewski and Weron, 2018; Ziel and Weron, 2018).

3.3.3. LASSO QRA

Regularization applied to quantile regression is not a completely new idea (for a mathematical treatment of this topic see Li and Zhu, 2008). There are even a few forecasting papers where this approach has been considered. On one hand, there are publications where LASSO is combined with QR to construct probabilistic forecasts of economic variables (Manzan, 2015), global radiation (Ben Bouallègue, 2017), electric demand (Lebotsa et al., 2018; He et al., 2019) or photovoltaic power (Agoua et al., 2019), for a pool of explanatory variables (but not point forecasts as in QRA). On the other, there are studies where penalized (via LASSO) QR is used to obtain a quantile of the predictive distribution (e.g., Value at Risk at the 95% level) from a pool of forecasts of this particular quantile (Bayer, 2018; Bracale et al., 2019). However, to our best knowledge, there is no paper that combines regularization with QRA, i.e., with point (not quantile) forecasts taken as independent variables in Eqn. (4).

The LASSO QRA or LQRA for short is constructed by applying the check function defined in Eqn. (5) to the ℓ^1 regularization in Eqn. (7):

$$\hat{\beta}_\alpha = \underset{\beta_\alpha}{\operatorname{argmin}} \left\{ \underbrace{\sum_{d,h} \left(\alpha - \mathbb{1}_{P_{d,h} < X_{d,h} \beta_\alpha} \right) \left(P_{d,h} - \sum_{n=1}^N \beta_\alpha^{(n)} \hat{P}_{d,h}^{(n)} \right)}_{\text{check function, see Eqn. (5)}} + \underbrace{\lambda \sum_{n=1}^N |\beta_\alpha^{(n)}|}_{\text{LASSO penalty}} \right\}, \quad (8)$$

where, as in Eqn. (4), $X_{d,h}$ is the matrix of independent variables, i.e., $\hat{P}_{d,h}^{(n)}$ for $n = 1, \dots, N$. Note, that Eqn. (8) can be rewritten to fit into the definition of a linear problem, thus to estimate $\hat{\beta}_\alpha$ we will use a basic linear programming tool, i.e., the Simplex algorithm (Dantzig et al., 1955).

Due to the linear penalty factor, LQRA selects the most informative point forecasts, so there is no need for experts to select them *a priori*. What is more, because of regularization, the final forecasts of different quantiles may be constructed using different sets of point forecasts. Potentially, this may improve the predictive performance. Actually, we expect the model to be at least as accurate as QRA and at the same time much more robust to a large number of point forecast (inputs). The higher the λ , the fewer point forecasts will finally be used to construct the quantile forecasts. Note, that in this study λ will not depend on α , i.e., we will present results for models with one λ for all 99 percentiles. In future research, it may be beneficial to test whether allowing λ to depend on α improves the forecasting accuracy.

To check the performance of LQRA we will conduct empirical tests on a logarithmic grid of 19 λ 's ranging from 10^{-1} to 10^3 . We will additionally include $\lambda = 0$, which corresponds to the original QRA method; in total we will report results for 20 different values of λ . Note, that in this paper we will not consider the optimal selection of the tuning parameter.

3.3.4. Models

The probabilistic forecasts are obtained for QR-type models calibrated on a rolling window of 364 days; this is in contrast to point forecasts, which are fitted on 701 windows of lengths ranging from 28 to 728 days. The initial calibration window covers the period from 29 September 2016 to 27 September 2017 and yields forecasts for all 24 hours of the next day (28 September). Afterwards, the window is rolled forward by one day and the models are re-estimated to obtain forecasts for 29 September. This procedure is repeated until the predictions for 31 December 2018 are made.

To obtain probabilistic forecasts of the day-ahead price, we will use only 25 out of 701 individual point forecasts available (see Section 4.1). Marcjasz et al. (2018) claim that it is not beneficial to take into consideration models that are calibrated on windows of fewer than 56 days. What is more, forecasts obtained for windows of similar lengths provide almost identical results. Thus, as the potential variables for LQRA we have selected point forecasts obtained for models calibrated on windows of 56, 84, 112, $\dots$, 728 days; see the black dots in Figure 3. The resulting models are denoted by **LQRA**(λ_i), where $\lambda_i = 10^{\frac{2(i-1)}{9}-1}$ for $i = 1, \dots, 19$ is a logarithmic grid of λ 's ranging from 10^{-1} to 10^3 .

Note, that $\lambda = 0$ in Eqn. (8) corresponds to 'pure' or non-regularized QRA. The latter is treated as the first benchmark and denoted later in the text by **QRA**. The next 25 benchmarks use QR applied to individual point forecasts; we denote them by **QR**(T), where $T = 56, 84, 112, \dots, 728$. The last three benchmarks are motivated by the results of Marcjasz et al. (2020). The authors notice that averaging forecasts can significantly improve the accuracy. Thus, we additionally average probabilistic forecasts **QR**(T), both 'horizontally' and 'vertically', see Section 3.3.2 in the cited article. We denote them by **Q-Ave** and **F-Ave**, respectively. Finally, since it may be beneficial to first average point forecasts and then apply QR (instead of averaging predictions on the probabilistic level), we also consider the Quantile Regression Machine (QRM) approach of Uniejewski et al. (2019). In this model, denoted by **QRM**, all 25 individual point forecasts are first averaged and the result is then used as an input to the QR algorithm. Interestingly, QRM yields nearly identical forecasts to the Factor QRA (or FQRA) model of Maciejowska et al. (2016), because the latter applies QR to the first factor (which approximates the mean) extracted from a pool of forecasts.

Finally, note that in this study we forecast 99 percentiles, which can be treated as a relatively good approximation of the entire distribution. Due to a possible lack of monotonicity of the quantile forecasts, after obtaining predictions of the 99 percentiles we sort the results, independently for each day and hour.

Figure 3: The *Mean Absolute Errors* (MAE) for the POLEX (*left*) and Nord Pool (*right*) datasets from the period 29 September 2016 to 31 December 2018 as a function of the calibration window length ranging from 28 to 728 days. Black dots represent forecasts used as inputs for QR-based procedures, gray dots – predictions for all considered window lengths.

4. Results

4.1. Point forecasts

The point forecasts are evaluated using the *Mean Absolute Error*:

$$\text{MAE} = \frac{1}{24D} \sum_{d=1}^D \sum_{h=1}^{24} |\hat{P}_{d,h} - P_{d,h}|,$$

where D is the number of days in the out-of-sample test period. The results are presented in Figure 3. They are very similar to those reported in Hubicka et al. (2019) and Marcjasz et al. (2018). As can be seen for both datasets, the optimal (in terms of MAE) calibration window length is around 100-days; this can hardly be selected *ex-ante*. For both datasets the shortest windows are not recommended. What was also reported in Marcjasz et al. (2018), for some datasets like Nord Pool, the longer calibration window we take, the worse the autoregressive model performs. For the POLEX market we do not see such a behavior.

4.2. Probabilistic forecasts

4.2.1. Error measures and testing for predictive ability

When evaluating probabilistic forecasts, we follow the paradigm of ‘maximizing sharpness subject to reliability’ (Gneiting and Katzfuss, 2014; Nowotarski and Weron, 2018). To check the *reliability* we compute the empirical coverage:

$$I_{d,h}^{\alpha} = \begin{cases} 1, & \text{for } P_{d,h} \in [\hat{L}_{d,h}^{\alpha}, \hat{U}_{d,h}^{\alpha}], \\ 0, & \text{for } P_{d,h} \notin [\hat{L}_{d,h}^{\alpha}, \hat{U}_{d,h}^{\alpha}], \end{cases} \quad (9)$$

where $\hat{L}_{d,h}^{\alpha}$ is the lower and $\hat{U}_{d,h}^{\alpha}$ the upper bound of the prediction interval (PI), and compare it to the nominal coverage. To this end, we perform the Kupiec (1995) test for the 50% and 90% PIs.

Figure 4: Empirical coverage for the POLEX (*top*) and Nord Pool (*bottom*) dataset from the period 28 September 2017 to 31 December 2018. The black dashed line corresponds to the expected coverage of 90%. Results are divided into two plots, the coverage for benchmark models (*left*; see Section 3.3.4) and **LQRA** as a function of λ (*right*; the filled blue circles represent the three selected values of λ considered in Sections 4.2.3 and 4.2.5).

Then, we use the Pinball Score (or Pinball Loss) to evaluate the *sharpness*. It is a proper scoring rule and a special case of an asymmetric piecewise linear loss function (Gneiting, 2011):

$$\text{PS}(\hat{q}_{\alpha,P}, P_{d,h}, \alpha) = \begin{cases} (1 - \alpha)(\hat{q}_{\alpha,P} - P_{d,h}) & \text{for } P_{d,h} < \hat{q}_{\alpha,P}, \\ \alpha(P_{d,h} - \hat{q}_{\alpha,P}) & \text{for } P_{d,h} \geq \hat{q}_{\alpha,P}, \end{cases} \quad (10)$$

where $\hat{q}_{\alpha,P}$ is the price quantile of order $\alpha \in (0, 1)$ and $P_{d,h}$ is the observed price for day d and hour h . The lower PS is, the more accurate are the probabilistic forecasts for a given quantile. The Pinball Score can be averaged across all percentiles (i.e., $\alpha = 0.01, 0.02, \dots, 0.99$; as in the GEFCom2014 competition, see Hong et al., 2016) and all hours in the whole out-of-sample test period. This yields the *Aggregate Pinball Score* (APS), which is equivalent to computing the quantile representation of the so-called Continuous Ranked Probability Score (CRPS), see Laio and Tamea (2007).

Finally, to draw statistically significant conclusions on the outperformance of forecasts of one model by those of another, we use the Giacomini and White (2006) test for *conditional predictive ability* (CPA). It can be regarded as a generalization of the commonly used Diebold-Mariano test for *unconditional* predictive ability. Here, one statistic for each pair of models is computed based

on the 24-dimensional vector of Pinball Scores for each day:

$$\Delta_{X,Y,d} = \|\text{APS}_{X,d}\| - \|\text{APS}_{Y,d}\|,$$

where $\|\text{APS}_{X,d}\| = \sum_{h=1}^{24} \sum_{\alpha=0.01}^{0.99} \text{PS}(\hat{q}_{\alpha,P}, P_{d,h}, \alpha)$ for model X . For each pair of models and both datasets we compute the p -values of the CPA test with null $H_0 : \phi = 0$ in the regression: $\Delta_{X,Y,d} = \phi' \mathbb{X}_{d-1} + \varepsilon_d$, where $\mathbb{X}_{d-1}$ contains information of day $d - 1$, i.e., a constant and lags of $\Delta_{X,Y,d}$.

4.2.2. Empirical coverage

In Figure 4 we depict the empirical coverage for the 90% PI constructed from the 5th and 95th percentiles, i.e., for $\alpha = 0.05$ and 0.95 . Plotted are the results of empirical coverage aggregated for all hours from whole out-of-sample period (28 September 2017 to 31 December 2018; $460 \times 24 = 11\,040$ observations). Several conclusions can be drawn:

- Firstly, we can observe that the dashed lines indicating the expected level of 90% are never reached. This means that all PIs are too narrow.
- As far as benchmarks are concerned, the best performing is **F-Ave**, but also **Q-Ave** beats all **QR(T)** models. This indicates that averaging probabilistic forecasts improves their reliability. The outperformance of ‘horizontal’ by ‘vertical’ averaging supports the results of Uniejewski et al. (2019).
- **LQRA** strongly depends on λ . Rather larger values of the tuning parameter are preferable.
- The most reliable results are obtained with **F-Ave** for POLEX and **LQRA** with $\lambda_{18} = 599.48$ (≈ 599) for Nord Pool. They yield a coverage of 89.5% and 88.0%, respectively.

Similar conclusions can be drawn for the 50% PIs (not plotted here for the sake of clarity).

4.2.3. The Kupiec test

The test is performed to check whether the obtained PIs are close enough to the nominal values of 50% or 90%. We report the results only for the three best performing benchmarks (**Q-Ave**, **F-Ave**, **QRM**) and **LQRA** with three well performing values of the tuning parameter: $\lambda_{10} = 10$, $\lambda_{14} = 77.43$ (≈ 77) and $\lambda_{18} = 599.48$ (≈ 599), denoted by **LQRA(10)**, **LQRA(77)** and **LQRA(599)**, respectively, and indicated by filled blue circles in Figures 4 and 6. The Kupiec (1995) test is conducted for the whole out-of-sample period but separately for each of the 24 hourly series. Based on the likelihood ratio statistics depicted in Figure 5 we can draw the following conclusions:

- In the night hours, all considered benchmarks struggle to pass the test for the 90% PI at any significance level. For both datasets, the Kupiec statistics for **Q-Ave**, **F-Ave** and **QRM** is close to exceeding or exceeds 20 for hours 1-6 and 24. For the remaining hours it is hard to tell whether **Q-Ave** or **QRM** is better.
- **LQRA** outperforms the benchmarks. The Kupiec statistic exceeds 20 for only a few hours and night hours are no longer a problem. Most of the hours pass the test at the 5% significance level.

Figure 5: The likelihood ratio (LR) statistics of the Kupiec test for 50% and 90% PIs computed for three selected benchmarks and LQRA with three well performing values of λ (see text for details) for the POLEX (*top 6 panels*) and Nord Pool (*bottom 6 panels*) datasets. The test was performed separately for each hour of the day. The dashed (dotted) horizontal lines refer to the 5% (1%) significance level. All values exceeding 20 are capped at this threshold.

Figure 6: The Aggregate Pinball Score (APS) for the POLEX (*top*) and Nord Pool (*bottom*) datasets for the period 28 September 2017 to 31 December 2018, plotted separately for benchmark models (*left*; see Section 3.3.4) and **LQRA** as a function of λ (*right*; the filled blue circles represent the three selected values of λ considered in Sections 4.2.3 and 4.2.5).

- Again we can see how different are the results obtained for different values of λ . For POLEX, **LQRA(77)** and **LQRA(599)** are comparable and outperform **LQRA(10)** by a considerable margin for most of the hours. For the Nord Pool market, the differences across λ 's are smaller, but we can clearly see that **LQRA(599)** outperforms the other two.
- The Kupiec test indicates that the most reliable method is **LQRA(599)**, however, **LQRA(77)** performs comparably for the POLEX dataset.

4.2.4. Sharpness

In Figure 6 we depict the Aggregate Pinball Score (APS), i.e., the PS averaged across all 99 percentiles and all days and hours in the out-of-sample period. Like in Section 4.2.2, we present the results for all benchmarks and all considered values of λ for **LQRA**. We can observe that:

- For the Polish market the results are not much different from those presented in Figure 4. Again, the best performing benchmark is **F-Ave** with an APS of 7.05.
- For the POLEX dataset, all results obtained for **LQRA** with $\lambda > 10$ outperform the benchmarks by a large margin. The APS as function of λ has a ‘minimum’ at 6.81, ca. 3.4% lower compared to the APS of the best benchmark.

Figure 7: Results of the conditional predictive ability (CPA) test of Giacomini and White (2006) for forecasts of selected models for the POLEX (*left*) and Nord Pool (*right*) datasets. We use a heat map to indicate the range of the p -values – the closer they are to zero ($\rightarrow$ dark green) the more significant is the difference between the forecasts of a model on the X -axis (better) and the forecasts of a model on the Y -axis (worse).

- For the Nord Pool dataset the best performing benchmark is **QR(84)** with an APS of ca. 0.92. Note, that the APS results contradict the conclusions drawn from analyzing the empirical coverage in Section 4.2.2.
- For the Nordic market we can also conclude that most of the forecasts obtained for **LQRA** outperform the benchmarks by a large margin. The APS as a function of λ takes its ‘minimum’ at 0.88, more than 4% lower compared to the best benchmark.
- For both datasets **LQRA** as a function of λ is slowly varying close to the minimum. This suggests that the ‘close to optimal’ tuning parameter can be selected using expert knowledge.

4.2.5. Testing conditional predictive ability

As in Section 4.2.3, we perform the CPA test of Giacomini and White (2006) only for the three well performing benchmarks (**Q-Ave**, **F-Ave**, **QRM**) and **LQRA** with three well performing values of λ . In Figure 7 we illustrate the obtained p -values using ‘chessboards’ (analogously as in Serafin et al., 2019), i.e., we use a heat map to indicate the range of the p -values – the closer they are to zero ($\rightarrow$ dark green) the more significant is the difference between the forecasts of a model on the X -axis (better) and the forecasts of a model on the Y -axis (worse). Evidently, the CPA test results confirm and emphasize the observations made in Section 4.2.4. In particular, **F-Ave** significantly outperforms all other benchmarks and for both datasets the best performing model overall is **LQRA(77)**. Interestingly, even though both datasets are very different in behavior we can select one value of the tuning parameter, i.e., $\lambda_{14} = 77.43$, that yields an LQRA model that is never significantly outperformed by any other method.

5. Conclusions

In this paper we have introduced a new technique for computing probabilistic forecasts – LASSO QRA or LQRA for short. It is a regularized variant of Quantile Regression Averaging, which utilizes the Least Absolute Shrinkage and Selection Operator (LASSO) to select the valuable and eliminate the redundant point forecasts used as inputs to quantile regression. Although illustrated on datasets comprising day-ahead electricity prices, we believe that LQRA can be a useful approach also for other forecasting applications, in particular, in risk management.

We have evaluated the new technique using datasets from the Polish and Nordic power markets, a set of 25 point forecasts obtained for calibration windows of different lengths and 20 different values of the regularization parameter λ . By comparing against nearly 30 benchmarks, we provide evidence for the superior predictive performance of LQRA in terms of the Kupiec test for (unconditional) coverage, the pinball score and the test for conditional predictive accuracy (CPA). The problem of the optimal choice of λ , possibly changing across time (as in Uniejewski and Weron, 2018) and quantiles, a more thorough prediction error analysis (e.g., performance in the center vs. in the tails of the distribution, as in Janke and Steinke, 2019) or assessing the impact of different fundamental variables on the obtained price forecasts (see, e.g., Maciejowska, 2020) are left for future research.

Acknowledgments

This work was partially supported by the Ministry of Science and Higher Education (MNiSW, Poland) through Diamond Grant No. 0199/DIA/2019/48 (to BU) and the National Science Center (NCN, Poland) through MAESTRO grant No. 2018/30/A/HS4/00444 (to RW).

References

- Agoua, X., Girard, R., Kariniotakis, G., 2019. Probabilistic models for spatio-temporal photovoltaic power forecasting. *IEEE Transactions on Sustainable Energy* 10 (2), 780–789.
- Bayer, S., 2018. Combining Value-at-Risk forecasts using penalized quantile regressions. *Econometrics and Statistics* 8, 56–77.
- Ben Bouallègue, Z., 2017. Statistical postprocessing of ensemble global radiation forecasts with penalized quantile regression. *Meteorologische Zeitschrift* 26 (3), 253–264.
- Bracale, A., Carpinelli, G., Falco, P. D., 2019. Developing and comparing different strategies for combining probabilistic photovoltaic power forecasts in an ensemble method. *Energies* 12 (6), 1–16.
- Bunn, D., Andresen, A., Chen, D., Westgaard, S., 2016. Analysis and forecasting of electricity price risks with quantile factor models. *Energy Journal* 37 (1), 101–122.
- Dantzig, G. B., Orden, A., Wolfe, P., et al., 1955. The generalized simplex method for minimizing a linear form under linear inequality restraints. *Pacific Journal of Mathematics* 5 (2), 183–195.
- Gaillard, P., Goude, Y., Nedellec, R., 2016. Additive models and robust aggregation for GEFCom2014 probabilistic electric load and electricity price forecasting. *International Journal of Forecasting* 32 (3), 1038–1050.
- Giacomini, R., White, H., 2006. Tests of conditional predictive ability. *Econometrica* 74 (6), 1545–1578.
- Gneiting, T., 2011. Quantiles as optimal point forecasts. *International Journal of Forecasting* 27 (2), 197–207.
- Gneiting, T., Katzfuss, M., 2014. Probabilistic forecasting. *Annual Review of Statistics and Its Application* 1, 125–151.
- Hagfors, L., Bunn, D., Kristoffersen, E., Staver, T., Westgaard, S., 2016. Modeling the UK electricity price distributions using quantile regression. *Energy* 102, 231–243.

- He, Y., Qin, Y., Lei, X., Feng, N., 2019. A study on short-term power load probability density forecasting considering wind power effects. *International Journal of Electrical Power and Energy Systems* 113, 502–514.
- Hong, T., Pinson, P., Fan, S., Zareipour, H., Troccoli, A., Hyndman, R. J., 2016. Probabilistic energy forecasting: Global Energy Forecasting Competition 2014 and beyond. *International Journal of Forecasting* 32 (3), 896–913.
- Hubicka, K., Marcjasz, G., Weron, R., 2019. A note on averaging day-ahead electricity price forecasts across calibration windows. *IEEE Transactions on Sustainable Energy* 10 (1), 321–323.
- Janke, T., Steinke, F., 2019. Forecasting the price distribution of continuous intraday electricity trading. *Energies* 12 (22), 4262.
- Koenker, R. W., 2005. *Quantile Regression*. Cambridge University Press.
- Kostrzewski, M., Kostrzewska, J., 2019. Probabilistic electricity price forecasting with Bayesian stochastic volatility models. *Energy Economics* 80, 610–620.
- Kupiec, P. H., 1995. Techniques for verifying the accuracy of risk measurement models. *The Journal of Derivatives* 3 (2), 73–84.
- Laio, F., Tamea, S., 2007. Verification tools for probabilistic forecasts of continuous hydrological variables. *Hydrology and Earth System Sciences Discussions* 11 (4), 1267–1277.
- Lebotsa, M. E., Sigauke, C., Bere, A., Fildes, R., Boylan, J. E., 2018. Short term electricity demand forecasting using partially linear additive quantile regression with an application to the unit commitment problem. *Applied Energy* 222, 104 – 118.
- Li, Y., Zhu, J., 2008. L1-norm quantile regression. *Journal of Computational and Graphical Statistics* 17, 163–185.
- Liu, B., Nowotarski, J., Hong, T., Weron, R., 2017. Probabilistic load forecasting via Quantile Regression Averaging on sister forecasts. *IEEE Transactions on Smart Grid* 8, 730–737.
- Maciejowska, K., 2020. Assessing the impact of renewable energy sources on the electricity price level and variability – a quantile regression approach. *Energy Economics* 85, 104532.
- Maciejowska, K., Nowotarski, J., 2016. A hybrid model for GEFCom2014 probabilistic electricity price forecasting. *International Journal of Forecasting* 32 (3), 1051–1056.
- Maciejowska, K., Nowotarski, J., Weron, R., 2016. Probabilistic forecasting of electricity spot prices using Factor Quantile Regression Averaging. *International Journal of Forecasting* 32 (3), 957–965.
- Manzan, S., 2015. Forecasting the distribution of economic variables in a data-rich environment. *Journal of Business & Economic Statistics* 33 (1), 144–164.
- Marcjasz, G., Serafin, T., Weron, R., 2018. Selection of calibration windows for day-ahead electricity price forecasting. *Energies* 11, 2364.
- Marcjasz, G., Uniejewski, B., Weron, R., 2020. Probabilistic electricity price forecasting with NARX networks: Combine point or probabilistic forecasts? *International Journal of Forecasting* (DOI: 10.1016/j.ijforecast.2019.07.002).
- Misioerek, A., Trück, S., Weron, R., 2006. Point and interval forecasting of spot electricity prices: Linear vs. non-linear time series models. *Studies in Nonlinear Dynamics & Econometrics* 10 (3), Article 2.
- Mpfumali, P., Sigauke, C., Bere, A., Mulaudzi, S., 2019. Day ahead hourly global horizontal irradiance forecasting – application to South African data. *Energies* 12 (18), 3569.
- Narajewski, M., Ziel, F., 2019. Econometric modelling and forecasting of intraday electricity prices. *Journal of Commodity Markets* (DOI: 10.1016/j.jcomm.2019.100107).
- Nowotarski, J., Weron, R., 2015. Computing electricity spot price prediction intervals using quantile regression and forecast averaging. *Computational Statistics* 30 (3), 791–803.
- Nowotarski, J., Weron, R., 2018. Recent advances in electricity price forecasting: A review of probabilistic forecasting. *Renewable and Sustainable Energy Reviews* 81, 1548–1568.
- Serafin, T., Uniejewski, B., Weron, R., 2019. Averaging predictive distributions across calibration windows for day-ahead electricity price forecasting. *Energies* 12 (13), 256.
- Tibshirani, R., 1996. Regression shrinkage and selection via the lasso. *Journal of the Royal Statistical Society B* 58, 267–288.
- Tikhonov, A. N., 1963. Solution of incorrectly formulated problems and the regularization method. *Soviet Mathematics Doklady* 4, 1035–1038.
- Uniejewski, B., Marcjasz, G., Weron, R., 2019. On the importance of the long-term seasonal component in day-ahead electricity price forecasting: Part II – Probabilistic forecasting. *Energy Economics* 79, 171–182.

- Uniejewski, B., Nowotarski, J., Weron, R., 2016. Automated variable selection and shrinkage for day-ahead electricity price forecasting. *Energies* 9, 621.
- Uniejewski, B., Weron, R., 2018. Efficient forecasting of electricity spot prices with expert and LASSO models. *Energies* 11, 2039.
- Uniejewski, B., Weron, R., Ziel, F., 2018. Variance stabilizing transformations for electricity spot price forecasting. *IEEE Transactions on Power Systems* 33, 2219–2229.
- Wang, Y., Zhang, N., Tan, Y., Hong, T., Kirschen, D., Kang, C., 2019. Combining probabilistic load forecasts. *IEEE Transactions on Smart Grid* 10 (4), 3664–3674.
- Zhang, W., Quan, H., Srinivasan, D., 2018. Parallel and reliable probabilistic load forecasting via quantile regression forest and quantile determination. *Energy* 160, 810–819.
- Zhang, Y., Liu, K., Qin, L., An, X., 2016. Deterministic and probabilistic interval prediction for short-term wind power generation based on variational mode decomposition and machine learning methods. *Energy Conversion and Management* 112, 208–219.
- Ziel, F., 2016. Forecasting electricity spot prices using LASSO: On capturing the autoregressive intraday structure. *IEEE Transactions on Power Systems* 31 (6), 4977–4987.
- Ziel, F., Weron, R., 2018. Day-ahead electricity price forecasting with high-dimensional structures: Univariate vs. multivariate modeling frameworks. *Energy Economics* 70, 396–420.

HSC Research Report Series 2019

For a complete list please visit <http://ideas.repec.org/s/wuu/wpaper.html>

- 01 *Electricity price forecasting* by Katarzyna Maciejowska and Rafał Weron
- 02 *Assessing the impact of renewable energy sources on the electricity price level and variability – a Quantile Regression approach* by Katarzyna Maciejowska
- 03 *Perspectives of smart meters' roll-out in India: an empirical analysis of consumers' awareness and preferences* by Yash Chawla, Anna Kowalska-Pyzalska and Anna Skowronska-Szmer
- 04 *Regularized Quantile Regression Averaging for probabilistic electricity price forecasting* by Bartosz Uniejewski and Rafał Weron